

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : C07D 215/22, 209/08, 263/58, 235/06, 471/04, A61K 31/44, 31/415, 31/42 // (C07D 471/04, 235:00, 221:00)	A1	(11) International Publication Number: <b>WO 96/12703</b>
		(43) International Publication Date: 2 May 1996 (02.05.96)

(21) International Application Number: PCT/US95/12263

(22) International Filing Date: 2 October 1995 (02.10.95)

(30) Priority Data:  
08/328,212 25 October 1994 (25.10.94) US

(60) Parent Application or Grant

(63) Related by Continuation

US

Filed on

08/328,212 (CON)

25 October 1994 (25.10.94)

(71) Applicant (for all designated States except US): G.D. SEARLE &amp; CO. [US/US]; Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): KHANNA, Ish, Kumar [IN/US]; 149 Brandywine Court, Vernon Hills, IL 60061 (US). STEALEY, Michael, Allan [US/US]; 502 Juniper Parkway, Libertyville, IL 60048 (US). WEIER, Richard, Mathias [US/US]; 240 Hickory Court, Lake Bluff, IL 60044 (US).


(74) Agents: HASTREITER, Roberta, L. et al.; G.D. Searle &amp; Co., Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US).

(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).

Published

With international search report.

(54) Title: HETEROALKYL AND HETEROARYLTHIOALKYL THIOPHENOLIC COMPOUNDS AS 5-LIPOXYGENASE INHIBITORS



## (57) Abstract

The present invention provides compounds of formula (I) and the pharmaceutically acceptable salts thereof, wherein:  $\text{R}^1$  and  $\text{R}^2$  are each alkyl;  $n$  is an integer of from 0 to 5;  $p$  is an integer of from 0 to 1;  $q$  is an integer of from 0 to 3;  $r$  is an integer of from 0 to 1; and  $\text{R}^3$  is a bicyclic group, with the proviso that when  $\text{R}^1$  and  $\text{R}^2$  are each alkyl, and  $p$ ,  $q$  and  $r$  are each 0,  $\text{R}^3$  is not a naphthylmethyl group. These compounds are inhibitors of 5-lipoxygenase, and are useful for the treatment of 5-lipoxygenase mediated conditions, including inflammation-associated disorders and allergic and hypersensitivity reactions. The present invention also provides pharmaceutical compositions comprising a therapeutically-effective amount of a compound of formula (I) in combination with a pharmaceutically-acceptable carrier, and methods for treating 5-lipoxygenase mediated conditions, inflammation-associated disorders and allergies in an animal comprising administering a therapeutically-effective amount of a compound of formula (I) to the animal.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

HETEROARALKYL AND HETEROARYLTHIOALKYL THIOPHENOLIC COMPOUNDS  
AS 5-LIPOXYGENASE INHIBITORS

Background of the Invention

5       1. Field of the Invention

The present invention relates to heteroaralkyl and heteroarylthioalkyl thiophenolic compounds. More particularly, the present invention relates to the novel compounds of Formula I, which inhibit 5-  
10 lipoxigenase, to pharmaceutical compositions containing one or more of these compounds in combination with a pharmaceutically-acceptable carrier, and to medical methods of treatment employing these compounds.

15       2. Background Information

It is well recognized that arachidonic acid, an essential unsaturated fatty acid, is enzymatically oxygenated to various products, including, prostaglandins, thromboxanes, the 5-, 11-, 12- and  
20 15-hydroxyeicosatetraenoic acids (HETEs, DIHETEs) and hydroperoxyeicosatetraenoic acids (HPETEs) and the leukotrienes, all of which have potent physiological effects. The leukotrienes, which are produced via the 5-lipoxygenase pathway, are major contributors to the  
25 onset of the symptoms of asthma, and mediators for immediate hypersensitivity reactions, inflammation and other allergic responses.

Leukotrienes are found in inflammatory exudates, and are involved in the process of cellular invasion during inflammation. The term "leukotrienes" is used  
30 as a generic term to describe a class of substances, such as slow-reacting substance (SRS) which is an important mediator in asthma and other hypersensitivity reactions. Immunologically generated SRS is usually  
35 referred to as slow-reacting substance of anaphylaxis (SRS-A). SRS-A consists of leukotrienes (LT) known as A<sub>4</sub>, B<sub>4</sub>, C<sub>4</sub>, D<sub>4</sub>, and E<sub>4</sub>. LTC<sub>4</sub> is at least 100 times more potent than histamine in causing long lasting

- 2 -

bronchoconstricting effects. The leukotrienes also increase vascular permeability and cause decreased cardiac output and impaired ventricular contraction. LTB<sub>4</sub> may be an important mediator of inflammation in,  
5 for example, inflammatory bowel disease.

Chemotaxis is a reaction by which the direction of migration of cells is determined by substances in their environment. It is one of the major processes bringing leukocytes from the blood to an inflammatory site,  
10 whether the inflammation is caused by an infectious agent, allergic challenge, or other pro-inflammatory stimuli. LTB<sub>4</sub> is not only chemotactic for neutrophils and monocytes, but is also highly active in stimulating eosinophil locomotion. LTB<sub>4</sub> also stimulates calcium  
15 influx and aggregation of polymorphonuclear leukocytes and LTB<sub>4</sub> may, thus, play an important role in mediating both acute and chronic inflammation.

Rheumatoid spondylitis is characterized by an acute neutrophil flareup in the joint which is  
20 associated with elevated levels of LTB<sub>4</sub>. LTB<sub>4</sub> is also present in gouty effusions, and exposure to urate crystals is known to stimulate LTB<sub>4</sub> production by neutrophils. Accordingly, the 5-lipoxygenase inhibitors of the present invention through inhibition  
25 of neutrophil attraction and activation in arthritic joints should reduce the protease and oxidative burden believed responsible for joint destruction in arthritic diseases.

Aspirin and the other non-steroidal  
30 anti-inflammatory agents (NSAIDs) such as indomethacin, ibuprofen, fenoprofen, and the like, inhibit the synthesis of prostaglandins via the cyclooxygenase pathway of arachidonic acid metabolism. These prostaglandin synthetase inhibitors generally exhibit  
35 anti-inflammatory, anti-pyretic and analgesic activity, and are widely used in the treatment of arthritis. The non-steroidal anti-inflammatory agents can lead to the

- 3 -

formation of additional pro-inflammatory derivatives of arachidonic acid produced through the 5-lipoxygenase pathway which play a role in immediate hypersensitivity reactions and also have pronounced inflammatory effects. Administration of the NSAIDs alone can produce allergic reactions including bronchospastic reactivity, skin rashes, syndrome of abdominal pain, fever, chills, nausea and vomiting, and anaphylaxis. For this reason, aspirin and the other non-steroidal anti-inflammatory agents (NSAIDs) are generally contraindicated for patients suffering from asthma, or who have previously exhibited allergic sensitivity to aspirin or other NSAIDs. Co-administration of the 5-lipoxygenase inhibitors of this invention with cyclooxygenase inhibitors may mitigate the untoward side effects of the latter and allow the increased advantageous use of such cyclooxygenase inhibitors.

Prior to the recognition of the significance of the 5-lipoxygenase pathway of arachidonic acid metabolism in allergic reactions and inflammation, the search for effective therapeutic agents was based primarily on those agents which treated the symptoms of allergy and inflammation. There has since been effort to develop new drugs which selectively block the formation of the mediators of these conditions, and the present invention provides new chemical entities which are inhibitors of the 5-lipoxygenase pathway, and are useful in the treatment of asthma, rheumatoid arthritis, psoriasis, and other allergic, hypersensitivity, and inflammatory conditions.

See Bengt Samuelsson, "Leukotrienes: Mediators of Immediate Hypersensitivity Reactions and Inflammation", Science, Vol. 220, pp. 568-575 (May 1983); Michael K. Bach, "Inhibitors of Leukotriene Synthesis and Action", The Leukotrienes, Chemistry and Biology, pp 163-194 (Academic Press, Inc., 1984); C. W. Lee et al., "Human Biology and Immunoreactivity of Leukotrienes", Advances

- 4 -

in Inflammation Research, Volume 6, pp 219-225 (Raven Press, New York 1984); Editorial, "Leukotrienes and other Lipoxygenase Products in the Pathogenesis and Therapy of Psoriasis and Dermatoses", Arch. Dermatol ,  
5 Vol. 119, pp 541-547 (July, 1983); Robert A. Lewis et al., "A Review of Recent Contributions on Biologically active Products of Arachidonate Conversion", Int. J. Immunopharmac. , Vol. 4, No. 2, pp 85-90 (1982);  
10 Michael K. Bach, Biochemical Pharmacology , Vol. 33, No. 4, pp 515-521 (1984); E. L. Becker, Chemotactic Factors of Inflammation , pp 223-225 (Elsevier Science Publishers B.V., Amsterdam, 1983); P. Sharon and W.F. Stenson, Gastroenterology , Vol. 86, 453-460 (1984);  
15 and Musch, M.W. et al., Science , Vol. 217, 1255 (1982), each of which is incorporated herein by reference.

U.S. Patent No. 4,692,532 discloses substituted N-(4-Hydroxyphenylthioalkyl)cyclic imides.

20 U.S. Patent No. 4,808,644 discloses substituted (4-Hydroxyphenylthioalkyl) derivatives.

European Patent Application Publication Number 0 434 405 A1 discloses imidazo[4,5-c]pyridines.

Japanese Patent Application No. 04001177 discloses the preparation of sulfonium compounds as hardening  
25 initiators.


Zotova et al., "Synthesis and Biological Activity of Substituted Sulfides of Indole and Benzofuran," Khim. - Farm. Zh., 26(1), 52-5, disclose the synthesis of substituted sulfides of indole and benzofuran.

30 Each of the documents described hereinabove discloses compounds which are structurally different from the compounds of the present invention. Thus, the compounds of the present invention are structurally distinct from that which has been described in the art.

- 5 -

SUMMARY OF THE INVENTION

The present invention provides compounds having a structure of Formula I:



Formula I

and the pharmaceutically acceptable salts thereof,  
wherein:

15 R<sup>1</sup> and R<sup>2</sup> are each alkyl;

n is an integer of from 1 to 5;

p is an integer of from 0 to 1;

20

q is an integer of from 0 to 3;

r is an integer of from 0 to 1; and

25 R<sup>3</sup> is a bicyclic group,

with the proviso that when R<sup>1</sup> and R<sup>2</sup> are each alkyl, and  
p, q and r are each 0, R<sup>3</sup> is not a naphthylmethyl group.

- 6 -

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS(1) Description of Invention

In one aspect, the present invention provides  
5 compounds comprising a structure of Formula I, as  
described above, and pharmaceutically-acceptable salts  
thereof.

Specific compounds within the scope of the  
invention include, but are not limited to, the  
10 compounds discussed in the examples presented below, as  
well as their pharmaceutically-acceptable salts.

Contemplated equivalents of the compounds  
described in Formula I include compounds which  
otherwise correspond thereto, and which have the same  
15 general properties thereof, wherein one or more simple  
variations of substituents are made which do not  
adversely affect the efficacy of the compound.

Certain compounds of the present invention may  
contain a basic functional group, such as amino and the  
20 like, and are, thus, capable of forming  
pharmaceutically-acceptable salts with  
pharmaceutically-acceptable acids. The term  
"pharmaceutically-acceptable salts" in this respect,  
refers to the relatively non-toxic, inorganic and  
25 organic acid addition salts of compounds of the present  
invention. These salts can be prepared in situ during  
the final isolation and purification of the compounds  
of the invention, or by separately reacting a purified  
compound of the invention in its free base form with a  
30 suitable organic or inorganic acid, and isolating the  
salt thus formed. Representative salts include the  
hydrobromide, hydrochloride, sulfate, bisulfate,  
phosphate, nitrate, acetate, valerate, oleate,  
palmitate, stearate, laurate, benzoate, lactate,  
35 phosphate, tosylate, citrate, maleate, fumarate,  
succinate, tartrate, naphthylate, mesylate,  
glucoheptonate, lactobionate, and laurylsulphonate



- 7 -

salts and the like. (See, for example, S.M. Berge et al., "Pharmaceutical Salts," J. Pharm. Sci., 66, 1-19 (1977), which, as well as all other documents referred to herein, is incorporated herein by reference.)

5 In other cases, the compounds of the invention may contain one or more acidic functional groups, such as carboxyl and the like, and, thus, are capable of forming pharmaceutically-acceptable salts with pharmaceutically-acceptable bases. The term  
10 "pharmaceutically-acceptable salts" in these instances refers to the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention. These salts can be prepared in situ during the final isolation and purification of the compounds,  
15 or by separately reacting the purified compound in its free acid form with a suitable base, such as the hydroxide, carbonate or bicarbonate of a metal cation, with ammonia, or with a pharmaceutically-acceptable organic primary, secondary or tertiary amine.  
20 Representative alkali or alkaline earth salts include the lithium, sodium, potassium, calcium, magnesium, and aluminum salts and the like. Representative organic amines useful for the formation of base addition salts include ethylamine, diethylamine, ethylenediamine,  
25 ethanolamine, diethanolamine, piperazine and the like. (See, for example, S. M. Berge et al., "Pharmaceutical Salts," J. Pharm. Sci., 66, 1-19 (1977), which is incorporated herein by reference).

In another aspect, the present invention provides  
30 pharmaceutically-acceptable compositions which comprise a therapeutically-effective amount of one or more of the compounds of Formula I, as described herein, formulated together with one or more pharmaceutically-acceptable carriers. The pharmaceutical compositions  
35 of the invention may be specially formulated for oral administration in solid or liquid form, or in the form

- 8 -

of an aerosol for inhalation, for parenteral injection, or for rectal or vaginal administration.

The present invention also comprises methods for treating 5-lipoxygenase mediated conditions, such as inflammation, or inflammation-associated disorders, and allergies in an animal, the method comprising administering to the animal having such condition a therapeutically-effective amount of a compound of Formula I.

10 The most preferred embodiment of the invention is the compound described in Example 14 below.

Certain compounds of this invention may exist in geometric or stereoisomeric forms. The present invention contemplates all such compounds, including cis- and trans-geometric isomers, R- and S-enantiomers, 15 diastereomers, d-isomers, l-isomers, the racemic mixtures thereof, and other mixtures thereof, as falling within the scope of the invention. Additional asymmetric carbon atoms may be present in a substituent 20 such as an alkyl group. All such isomers, as well as mixtures thereof, are intended to be included in this invention.

- 9 -

(2) Definitions

For purposes of clarity, the terms and phrases used throughout this specification and the appended claims are defined in the manner set forth directly below.

Some of the chemical structures which are presented in this specification and the appended claims have been drawn using the convention which employs lines to represent alkyl radicals, which is known by those of skill in the art.

The abbreviation "AIBN" as used herein means azobisisobutyronitrile.

The term "alkyl" as used herein defines straight or branched chain monovalent hydrocarbon radicals having between about 1 to about 10 carbon atoms, within which includes from about 1 to about 6 carbon atoms, and further within which includes from about 1 to about 3 carbon atoms. Representative alkyl radicals include, for example, methyl, ethyl, propyl, isopropyl, butyl, t-butyl, sec-butyl, isobutyl, pentyl, 1-methylbutyl, isopentyl, neopentyl, hexyl, octyl, nonyl, decyl, t-pentyl, etc.

The term "animal" as used herein includes mammals and nonmammals, and further includes humans and nonhuman mammals.

The term "bicyclic group" as used herein means a double ring structure comprising two 6-membered rings, two 5-membered rings or one 6-membered ring and one 5-membered ring (in any order), each of the two rings independently containing one, two or three double bonds, and the double ring structure containing 1, 2 or 3 nitrogen atoms and, optionally, containing further substitutions of up to 2 groups selected from oxygen,

$$\begin{array}{c} \text{O} \\ || \\ -\text{NH}, -\text{N-alkyl and } -\text{C-} \end{array}$$
 Representative bicyclic groups include 1(H)-imidazo[4,5-c]pyridine, 1(H)-imidazo[4,5-b]pyridine, benzimidazole, benzoxazole and the like.

- 10 -

The abbreviation "b.p." as used herein means boiling point.

The term "composition" as used herein means a product which results from the combining of more than one element or ingredient.

The abbreviation "DMA" as used herein means dimethylacetamide.

The abbreviation "DMF" as used herein means dimethylformamide.

The abbreviation "DMSO" as used herein means dimethylsulfoxide.

The abbreviation "DSC" as used herein means Differential Scanning Calorimetry.

The abbreviation "DTT" as used herein means dithiothreitol.

The abbreviation "EDTA" as used herein means ethylene diamine tetraacetic acid.

The abbreviation "EtOH" as used herein means ethanol.

The term "halogen" as used herein means chlorine, bromine, fluorine, and iodine.

The abbreviation "HETE" as used herein means hydroxyeicosatetraenoic acid.

The abbreviation "HPETE" as used herein means hydroperoxyeicosatetraenoic acid.

The term "heteroatom" as used herein means an atom of any element other than carbon or hydrogen.

The abbreviation "HPLC" as used herein means High Pressure Liquid Chromatography.

The abbreviation "ip" as used herein means intraperitoneal.

The abbreviation "MeOH" as used herein means methanol.

The abbreviation "mp" as used herein means melting point.

The abbreviation "mpk" as used herein means milligrams per kilogram.

- 11 -

The abbreviation "MPO" as used herein means myeloperoxidase.

The abbreviation "NMR" as used herein means Nuclear Magnetic Resonance.

5       The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, 10       intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection 15       and infusion.

      The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgement, suitable for use in 20       contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

      The phrase "pharmaceutically-acceptable carrier" 25       as used herein means a pharmaceutically-acceptable material, composition or vehicle, as defined directly above, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting a chemical compound or 30       pharmaceutical agent from one organ, or portion of the body, to another organ, or portion of the body. Some examples of materials which can serve as pharmaceutically-acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) 35       starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6)

- 12 -

gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laureate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) phosphate buffer solutions; and (21) other non-toxic compatible substances employed in pharmaceutical formulations.

The expression "pharmaceutically acceptable salts" is intended to include those salts capable of being formed with the compounds of the present invention without materially altering the covalent chemical structure thereof. Such salts include inorganic and organic base or acid addition salts, such as sodium, potassium, calcium, ammonium, alkylammonium, quaternary ammonium, triethanolamine, lysine, hydrochloride, hydrobromide, phosphate, citrate, etc. well known to those skilled in the art. The foregoing salts are prepared in the conventional manner by neutralization of the compounds of formula I with the desired base or acid.

The abbreviation "t-Bu" as used herein means tert-butyl.

The phrase "therapeutically-effective amount" as used herein means an amount of a compound, material, or composition which is an effective dose for eliminating or ameliorating pain in an animal, or for producing some other desired therapeutic effect, at a reasonable benefit/risk ratio applicable to any medical treatment.

The phrases "title compound," "title product" and "title material" as used herein mean that compound, product or material whose chemical name is given, and/or whose structure is shown, in the particular

- 13 -

example, or subpart thereof, referred to. If no particular example, or subpart thereof, is referred to, it means that compound, product or material whose chemical name is given, and/or whose structure is  
5 shown, in the particular example, or subpart thereof, in which it appears.

The abbreviation "TPA" as used herein means tetradecanoylphorbol 13-acetate.

- 14 -

(3) Utility

The present invention provides compounds which block the 5-lipoxygenase metabolic pathway and, therefore, block the formation of the leukotrienes responsible for allergy and inflammation, and represent therapeutic agents which are useful in the treatment of allergic and hypersensitivity reactions and inflammation, alone, or also may be utilized in combination with other lipoxygenase inhibitors or with cyclooxygenase inhibitors, such as the non-steroidal anti-inflammatory agents.

The compounds of Formula I which inhibit 5-lipoxygenase are useful in treating 5-lipoxygenase mediated conditions, such as inflammation and inflammation-associated disorders and allergies and related disorders and conditions, such as arthritis, asthma, and psoriasis.

Compounds of the present invention would be useful for the treatment of inflammation in an animal, and for treatment of other inflammation-associated disorders, such as an analgesic in the treatment of pain and headaches, or as an antipyretic for the treatment of fever. For example, compounds of the present invention would be useful to treat arthritis, including but not limited to rheumatoid arthritis, spondyloarthropathies, gouty arthritis, systemic lupus erythematosus, osteoarthritis and juvenile arthritis. Such compounds would be useful in the treatment of asthma, bronchitis, menstrual cramps, tendinitis, bursitis, and skin related conditions such as psoriasis, eczema, burns and dermatitis. Compounds of the present invention also would be useful to treat gastrointestinal conditions such as inflammatory bowel syndrome, Crohn's disease, gastritis, irritable bowel syndrome and ulcerative colitis. Compounds of the present invention would be useful in treating inflammation in such diseases as vascular diseases, migraine headaches, periarteritis nodosa, thyroiditis, aplastic anemia, Hodgkin's



- 15 -

disease, sclerodoma, rheumatic fever, type I diabetes, myasthenia gravis, sarcoidosis, nephrotic syndrome, Behcet's syndrome, polymyositis, hypersensitivity, conjunctivitis, gingivitis, swelling occurring after  
5 injury, myocardial ischemia, and the like. Compounds of the present invention are useful as anti-inflammatory agents, such as for the treatment of arthritis, with the additional benefit of having significantly less harmful side effects.

10

- 16 -

(4) Dosage and Mode of Administration

The compounds of the present invention can be administered to a patient in such oral dosage forms as tablets, capsules, pills, powders, granules, elixirs, or syrups as well as aerosols for inhalation. Likewise, administration may be effected intravascularly, subcutaneously, topically, or intramuscularly using dosage forms known to those of ordinary skill in the pharmaceutical arts. In general, the preferred form of administration is oral. An effective but non-toxic amount of the compound is employed in treatment. The dosage regimen utilizing the present compounds is selected in accordance with a variety of factors including the type, age, weight, sex, and medical condition of the patient, the severity of the condition to be ameliorated, and the route of administration. A physician of ordinary skill can readily determine and prescribe the effective amount of the drug required to prevent, treat or arrest the progress of the condition.

Dosages of the compounds of the present invention, will range generally between 0.1 mg/kg/day to about 100 mg/kg/day, and preferably between about 0.5 mg/kg of body weight per day to about 50 mg/kg of body weight per day, when administered to patients suffering from inflammation or allergic or hypersensitivity reactions. In general, a unit dose form of the compounds of the invention will contain from about 1.75 to about 750 mg of compound. The compound may be administered in divided dosages, e.g. two or more times daily. The compounds may also be administered transdermally or topically to treat proliferative skin conditions such as psoriasis. The daily dosage may be administered in a single dose or in equal divided doses three or four times daily.

Unit dosage forms such as tablets and capsules can contain any suitable, predetermined, therapeutically

- 17 -

effective amount of one or more active agent and a pharmaceutically acceptable carrier or diluent.

Generally speaking, solid oral unit dosage forms and other unit dosage forms of the compounds of this

5 invention will contain from 1.75 to 750 mg per tablet of drug as the effective lipxygenase inhibiting amount of the compound.

In the case of acute allergic or hypersensitivity reactions, it is generally preferable to administer the  
10 initial dosage via the parenteral route and continue parenteral administration until the patient is stabilized, and can be maintained, if necessary, on oral dosing.

In the case of psoriasis and other skin  
15 conditions, it may be preferred to apply a topical preparation of a compound of this invention to the affected area three or four times daily.

In treating asthma and arthritis with a compound of this invention, the compounds may be administered  
20 either on a chronic basis, or as symptoms appear. However, in the case of arthritis and other inflammatory conditions which can lead to deterioration of joints and malformations, it is generally preferable to administer the active agent on a chronic basis.

25 A typical tablet of this invention can have the following compositions:

	Ingredient	Mg/tablet
30	Active ingredient	100
	Starch, U.S.P.	57
	Lactose, U.S.P.	73
	Talc, U.S.P.	9
35	Stearic acid	12

In the pharmaceutical compositions and methods of the present invention, at least one of the active compounds of the invention, or a pharmaceutically  
40 acceptable salt thereof, will typically be administered

- 18 -

in admixture with suitable pharmaceutical diluents, excipients, or carriers (collectively referred to herein as "carrier" materials) suitably selected with respect to the intended form of administration, that

5 is, oral tablets, capsules, elixirs, syrups, and the like, and consistent with conventional pharmaceutical practices. For instance, for oral administration in the form of tablets or capsules, the active drug component may be combined with any oral non-toxic

10 pharmaceutically acceptable inert carrier such as lactose, starch, sucrose, cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol and the like. For oral administration in liquid form, the active drug component may be combined

15 with any oral non-toxic pharmaceutically acceptable inert carrier such as ethanol and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated in the mixture. Suitable

20 binders include starch, gelatin, natural sugars, corn sweeteners, natural and synthetic gums such as acacia, sodium alginate, carboxymethylcellulose, polyethylene glycol, and waxes. Lubricants for use in these dosage forms include boric acid, sodium benzoate, sodium

25 acetate, sodium chloride, and the like. Disintegrators include, without limitation, starch, methylcellulose, agar, bentonite, guar gum, and the like.

- 19 -

(5) Methods of Preparation

In general, the compounds of the present invention may be prepared by the methods illustrated in the following general reaction schemes, or by modifications thereof, from readily available starting materials in a conventional manner. Unless otherwise specified, the various substituents of the compounds are defined in the same manner as they are defined hereinabove in Formula I.

10 If a particular enantiomer of a compound of the present invention is desired, it may be prepared by chiral synthesis, or by derivation with a chiral auxiliary, where the resulting diastereomeric mixture is separated and the auxiliary group cleaved to provide  
15 the pure desired enantiomers. Alternatively, where the molecule contains a basic functional group, such as amino, or an acidic functional group, such as carboxyl, diastereomeric salts are formed with an appropriate optically-active acid or base, followed by resolution  
20 of the diastereomers thus formed by fractional crystallization or chromatographic means well known in the art, and subsequent recovery of the pure enantiomers.

25 The compounds of the present invention may be synthesized by the procedures described in General Reaction Schemes 1, 2 and 3 presented hereinbelow.

- 20 -

General Reaction Schemes 1, 2 and 3 describe the general methods for synthesizing the compounds of the present invention wherein  $R^1$  and  $R^2$  are each alkyl,  $n$  is an integer of from 1 to 5,  $p$  is an integer of from 0 to 1,  $q$  is an integer of from 0 to 3,  $r$  is an integer of from 0 to 1 and  $R^3$  is a bicyclic group.

The compounds of Formula I (wherein  $p$  and  $q = 0$ ) can be synthesized in the manner shown in General Reaction Scheme 1. Substituted 4-hydroxythiophenols (II) are alkylated with, for example, 3-bromo-1-propanol, 2-bromoethanol or ethylene oxide to give the hydroxy compounds III. This alkylation reaction is carried out in the presence of a non-nucleophilic base, such as sodium carbonate, potassium carbonate, sodium bicarbonate or diisopropylethylamine, using a solvent such as acetone, dimethylformamide or dimethylacetamide. Compounds III can also be synthesized by reacting compounds II with haloalkyl esters, such as methyl 4-bromobutyrate or 4-bromobutyric acid, to give compounds IV, followed by reduction with reducing agents, such as lithium borohydride for an ester or borane-tetrahydrofuran for a carboxylic acid. The alcohols III are converted to bromide derivatives V using a triarylphosphine, such as triphenylphosphine, in combination with a bromine source, such as carbon tetrabromide. This reaction is preferably carried out in an ethereal solvent, such as diethyl ether, tetrahydrofuran or dioxane at 20-50°C. Alternatively, the alcohols III may be first converted to sulfonate esters by reaction with a suitable sulfonylating agent, such as p-toluenesulfonyl chloride, benzenesulfonyl chloride or trifluoromethanesulfonic anhydride, followed by treatment with a bromide salt, such as lithium bromide, sodium bromide or potassium bromide. The bromo compounds V are then reacted with the heterocycle to give the target compounds VI. When compounds VI are


- 21 -

reacted with 1(H)-imidazo[4,5-c]pyridine in the absence of an additional base, the product of alkylation at the pyridine nitrogen predominates.

Alternatively, as illustrated in General Reaction Scheme 2, a heterocyclic compound, such as indole, benzimidazole, 1(H)-imidazo[4,5-c]pyridine or 2-mercaptobenzoxazole, may be first alkylated with, for example, 1-iodo-3-chloropropane, to give compounds VII. This reaction is preferably carried out in a solvent such as dimethylformamide or dimethylacetamide using a base such as sodium hydride or potassium hydride. The intermediates VII are then alkylated with the substituted 4-hydroxythiophenols II using procedures as described for the first step of General Reaction Scheme 1 to give the desired compounds VI.


The compounds of Formula I can also be synthesized by following General Reaction Scheme 3. The substituted 4-hydroxythiophenols II are alkylated with an unsaturated alkyl halide or sulfonate ester, such as  $X(CH_2)_nCH=CH_2$  ( $X = Br, Cl, I, \text{ or } OSO_2R$ ;  $n = 1-5$ ;  $R = \text{phenyl, p-methyl phenyl, methyl or } -CF_3$ ) to give compounds VIII using conditions described for the first step of General Reaction Scheme 1. The epoxidation of compounds VIII using an oxidizing agent, such as m-chloroperoxybenzoic acid, followed by reaction with the appropriately substituted heterocycle gives the target compounds I. Alternatively, the appropriately substituted heterocycles may be reacted with the halohydrins X to give the target compounds I. The intermediates X can be synthesized by opening of epoxides IX with, for example, bromotrimethylsilane, hydrogen chloride or hydrogen bromide. In compounds of the present invention where  $n = 1$ , the halohydrins X can be directly synthesized by reaction of compounds II with epichlorohydrin.


- 22 -

GENERAL REACTION SCHEME 1



- 23 -

GENERAL REACTION SCHEME 2

GENERAL REACTION SCHEME 3

- 25 -

The conditions for carrying out the individual steps in each of the general reaction schemes presented above are conventional, well-known, and capable of wide variation.

- 5        Other methods known in the art can also be used to synthesize the compounds of the present invention.

- 26 -

(6) Biological Evaluations

(a) Recombinant 5-Lipoxygenase Inhibition Assay

The compounds of the present invention were  
5 evaluated with respect to 5-lipoxygenase inhibition  
according to the Recombinant 5-Lipoxygenase Inhibition  
Assay described below.

Preparation of Human Recombinant 5-Lipoxygenase

10 Human 5-lipoxygenase expressed in E. coli as a  
soluble protein was prepared in the manner described by  
M. Noguchi et al., "Expression of Human 5-Lipoxygenase  
cDNA in E. Coli," FEBS Letters, Vol. 249, No. 2, 267-  
270 (1989), which is incorporated herein by reference.  
15 The human 5-lipoxygenase was partially purified by ion  
exchange chromatography and stored in 50 mM Tris-HCl  
buffer (pH 8.1) containing 150 mM NaCl, 2 mM EDTA, 2 mM  
DTT, and 50% glycerol at -70°C before use.

Materials

20 The following solutions were prepared as  
indicated:  $\text{CaCl}_2/\text{ATP}$ , 12.5 mM  $\text{CaCl}_2$  and 5 mM ATP in  $\text{H}_2\text{O}$ ;  
arachidonic acid, 4 mM in ethanol; potassium phosphate  
buffer, 50 mM potassium phosphate and 0.1 mM EDTA, pH  
25 7.2. Test compounds were first prepared at 10 mM in  
DMSO and further diluted to desirable concentrations  
with DMSO.

Assay of 5-Lipoxygenase Activity

30 The assay was carried out in a quartz cuvette to  
which 15  $\mu\text{l}$   $\text{CaCl}_2/\text{ATP}$ , 7.5  $\mu\text{l}$  arachidonic acid, 7.5  $\mu\text{l}$   
ethanol or test compound, and 710  $\mu\text{l}$  potassium  
phosphate buffer, in that order, were added and mixed.  
The mixture was prewarmed to 25°C. The enzymatic  
35 reaction was started by adding 10  $\mu\text{l}$  of 5-lipoxygenase  
enzyme (~ 0.2 mg protein), and the formation of the  
reaction product 5-HPETE/5-HETE at 25°C for 8 minutes

- 27 -

was determined by measuring the increase in the absorbance at 240 nm using a Beckman DU-7 spectrophotometer.

5     The results for this assay are presented in Table 1 hereinbelow. The results for this assay are expressed in terms of activity with respect to 5-lipoxygenase inhibition as an  $IC_{50}$  value (50% inhibitory concentration).

10

- 28 -

(b) Tetradecanoylphorbol Acetate (TPA) Assay

The compounds of the present invention were evaluated with respect to 5-lipoxygenase inhibition according to the Tetradecanoylphorbol Acetate (TPA) Assay which is described in Rao et al., "Comparative Evaluation of Arachidonic Acid (AA) - And Tetradecanoylphorbol Acetate (TPA)-Induced Dermal Inflammation." Inflammation, Vol. 17, No. 6, 723-741 (1993), which is incorporated herein by reference.

AA (arachidonic acid) and TPA (12-O-tetradecanoylphorbol 13-acetate, a protein kinase C activator) are widely employed agents to induce cutaneous inflammation in experimental animals.

The initiation of inflammatory responses by metabolites of AA via CO (cyclooxygenase) and LO (lipoxygenase) pathways, and suppression of acute inflammatory responses by inhibitors of CO and LO have established a role for metabolites of AA in acute inflammation.

In this study, selective 5-lipoxygenase inhibitors, such as zileuton, MK-886, LTB<sub>4</sub> and LTD<sub>4</sub> receptor agonists were evaluated in acute inflammatory responses elicited by the topical application of TPA to the ears of mice. The increases in edema and myeloperoxidase (MPO) were used as readouts of inflammatory responses.

Animals. Female BALB/c mice (17-25 g; Jackson Labs, Bar Harbor, Maine) or male Swiss Webster mice (Charles River, Portage, Michigan) were used in the studies. The animals were housed in a vivarium with a 12-hour lighting schedule (0600-1800 hours on) with free access to food and water. Housing and experimental procedures were conducted according to institutionally approved (Monsanto, St. Louis, MO and

- 29 -

National Institute of Health) guidelines.

Drugs and Reagents. Hexadecyltrimethylammonium bromide (HTAB), and TPA were purchased from Sigma  
5 Chemical Co. (St. Louis, Missouri). Human polymorphonuclear leukocyte myeloperoxidase (20 units/mg protein) was obtained from Calbiochem (San Diego, California). One unit of MPO was defined as the activity of the enzyme that oxidizes 1  $\mu$ mol of hydrogen  
10 peroxide per minute at 25°C.

TPA Inflammation Model. Groups of mice (six to eight per treatment) received either vehicle (saline or 0.5% w/v methylcellulose or saline with 1% alcohol as  
15 appropriate) or drug intraperitoneally 60 minutes after the topical application of TPA [4  $\mu$ g/ear applied as a solution in acetone (200  $\mu$ g/ml), 10  $\mu$ l each to the inner and outer surface of ear]. The control group received 20  $\mu$ l of acetone as a topical application.  
20 Four hours following the application of TPA, increases in ear thickness were measured and ears were excised for MPO measurements.

Measurement of MPO. Ear samples were homogenized  
25 in a Brinkman polytron in 3 ml of 50 mM potassium phosphate buffer (pH 6.0), centrifuged (32,000 g, 20 minutes), and supernatants were discarded. Tissue pellets were reextracted into potassium phosphate buffer (50 mM, pH 6.0) with 0.5% HTAB, frozen-thawed

- 30 -

three times, and centrifuged to collect supernatants that were used in MPO assays. MPO assays were performed according to the procedures described by Bradley et al. [Bradley, P.B., et al. "Measurement of cutaneous inflammation: Estimation of neutrophil content with an enzyme marker," J. Invest. Dermatol. 78:206-209 (1982)], adapted to a 96-well plate format. Briefly, 7  $\mu$ l of unknowns or human neutrophil MPO standards were added to a 96-well plate. The reaction was initiated by the addition of 200  $\mu$ l of assay buffer containing 0.167 mg/ml o-dianisidine and 0.0005% hydrogen peroxide. The rate of change of absorbance was monitored in kinetic mode by a plate reader (Molecular Devices, Menlo Park, CA). Levels of MPO in lavage samples were determined from the calibration curves using human neutrophil MPO as the reference standard. The levels of MPO were expressed as milliunits per ear. Measurements of MPO, a granular constituent of neutrophils, were taken as indices of neutrophil influx into the inflamed tissue. MPO constitutes nearly 5% of neutrophil content and the utility of its measurements to correlate neutrophil influx is well established.

25        Data Analyses. Ear thickness measurements were averaged and responses in vehicle-treated groups were subtracted from those in other treatment groups to give



- 31 -

inflammatory responses. The percent inhibition was defined by the following equation:

$$\% \text{ inhibition} = \frac{(C_r - T_r)}{(C_r)} \times 100$$

where C<sub>r</sub> and T<sub>r</sub> refer to increases in ear thickness i.  
arachidonic acid- (or TPA-) treated and arachidonic  
5 acid- (or TPA-) + drug-treated groups, respectively.  
The data were analyzed by a one-way analysis of  
variance followed by a Neuman-Keuls or a parametric  
Dunnett's test, as known by those of skill in the art.

The results for this assay are presented in Table  
10 1 hereinbelow. The results for this assay are  
expressed in terms of activity with respect to 5-  
lipoxxygenase inhibition as a per cent inhibition value.  
A test compound which is active in this assay inhibits  
or prevents the increase in ear thickness induced by  
15 treatment with TPA.

- 32 -

(c) LTB<sub>4</sub> and Thromboxane Production by Calcium  
Ionophore Stimulated Human Blood - Human  
Whole Blood LTB<sub>4</sub> Assay

5 The compounds of the present invention were  
evaluated with respect to 5-lipoxygenase inhibition  
according to the Human Whole Blood LTB<sub>4</sub> Assay described  
below.

Human whole blood was collected in heparin  
containing Vacutainer brand tubes. The blood was  
10 diluted with RPMI-1640 cell culture media [1:4] (Gibco,  
Grand Island, NY) and 200 microliters were added per  
well in 96 well microtiter plates. Compounds of the  
present invention diluted in dimethylsulfoxide were  
added to the blood and allowed to incubate for 15  
15 minutes at 37°C in a humidified incubator. Calcium  
ionophore A23187 [20mcg/ml final concentration, Sigma  
Chemical Co., St. Louis, MO] was added, and the  
incubation was continued for 10 minutes. The  
incubation was terminated by centrifugation [833g, 10  
20 minutes at 4°C]. Supernatants were analyzed for LTB<sub>4</sub>  
and thromboxane by commercially available enzyme-linked  
immunoassays [Caymen Chemical Co., Ann Arbor, MI].  
Test compounds which inhibit 5-lipoxygenase in this  
assay inhibit the formation of LTB<sub>4</sub> and cause reduced  
25 levels of this agent in the supernatant. Test  
compounds which also inhibit cyclooxygenase in this  
assay also cause reduced levels of thromboxane in the  
supernatant.

The results for this assay are presented in Table  
30 1 hereinbelow. The results of this assay are expressed  
in terms of activity with respect to 5-lipoxygenase  
inhibition as an IC<sub>50</sub> value (50% inhibitory  
concentration).

- 33 -

TABLE 1

Compound Number	Example Number	Recombinant 5-Lipoxygenase Inhibition Assay IC <sub>50</sub> (μM)	TPA (Mouse, Ear) (50 mpk, ip + 120 minute) Edema MPO (% Inhibition)	Human Whole Blood Assay LTB <sub>4</sub> IC <sub>50</sub> (μM)	Human Whole Blood Assay Thromboxane IC <sub>50</sub> (μM)
5	3	0.57 ± 0.07	42	24	13.3
8	6	4.48 ± 0.36	*-	-	-
14	11	0.41 ± 0.02	-	-	-
15	12	0.49 ± 0.04	-	-	6.27
16	12	0.34 ± 0.10	-	-	14.4
18	13	0.32 ± 0.02	-	-	-
19	13	0.36 ± 0.07	-	-	-
21	14	0.14 ± 0.02	-	-	6.62
24	16	0.48 ± 0.09	-	-	2.26
26	17	0.93 ± 0.16	-	-	-
27	18	1.1 ± 0.23	46	-	11.1
29	20	1.8 ± 0.3	-	-	7.42
			-	-	20.1
			-	-	50

\*- = not calculated

- 34 -

(7) Examples

The following non-limiting examples further describe and illustrate details from the preparation of the compounds of the present invention from readily available starting materials. Those skilled in the art will readily understand and appreciate that known variations of the conditions and procedures in the following preparative methods can be utilized.

In these examples, all temperatures are degrees Celsius unless otherwise noted. Melting points were determined on a Fisher-Johns melting point apparatus or by DSC and are uncorrected.

Unless indicated otherwise in a particular example or biological evaluation, all of the starting materials, and all of the equipment, employed in the examples and biological evaluations are commercially available. Sources for these materials include Sigma Chemical Co. (St. Louis, MO), Aldrich Chemical Co. (Milwaukee, WI), Lancaster Synthesis (Windham, NH), Bioproducts For Science, Inc. (Indianapolis, IN), Fisher Scientific (Pittsburgh, PA), Boehringer Mannheim Biochemicals (Indianapolis, IN), Fluka Chemical Corp. (Ronkonkoma, NY), TCI, American Tokyo Kasei, Inc. (Atlanta, GA), Chemical Dynamics Corp. (South Plainfield, NJ), Amano International Enzyme Company, Inc. (Troy, VA), Nu-Chek-Prep, Inc. (Elysian, MN), Cayman Chemical (Ann Arbor, MI), Cistron (Pine Brook, NJ), Gibco/BRL (Gaithersburg, MD) and Invitrogen (Palo Alto, CA). Most of the starting materials were obtained from Aldrich Chemical Co. (Milwaukee, WI). The syntheses of those starting materials which are not commercially available are described in the examples presented below.


All patents and publications referred to in the examples, and throughout the specification, are hereby incorporated herein by reference, without admission that such is prior art.

- 35 -

Example 12,6-bis(1,1-dimethylethyl)-4-[(3-hydroxypropyl)thio]phenol (2)

5

10




To a solution of 2,6-bis(1,1-dimethyl)-4-mercaptophenol **1** (13.6 g, 57 mmol) in DMF (150 ml),  
15 anhydrous potassium carbonate (15.76 g, 114 mmol) and 3-bromopropanol (5.3 ml) were added. After stirring at room temperature for 20 hours, the reaction mixture was filtered and the filter cake was washed with methylene chloride. The combined filtrates were concentrated and  
20 the crude orange solid (14 g) obtained was chromatographed (silica gel; hexane/ethyl acetate 7/3) to give **2** (9.93 g, 59%) as a pale orange liquid. The structural assignment was supported by the <sup>1</sup>H-NMR spectrum in CDCl<sub>3</sub>.

- 36 -

Example 24-[(3-bromopropyl)thio]-2,6-bis(1,1-dimethylethyl)phenol (3)

5

10



Triphenylphosphine (17.54 g, 66.89 mmol) was added  
15 to a stirred solution of 2 (9.9 g, 33.44 mmol) and  
carbon tetrabromide (22.18 g, 66.89 mmol) in dry ether.  
After stirring at room temperature for 18 hours, the  
reaction mixture was concentrated and the crude product  
was chromatographed (silica gel; hexane/ethyl acetate  
20 98/2) to give 3 in quantitative yield. The structural  
assignment was supported by the <sup>1</sup>H-NMR spectra in CDCl<sub>3</sub>.


- 37 -

Example 3

2,6-bis(1,1-dimethylethyl)-4-[[3-(5H-imidazo-  
[4,5-c]pyridin-5-yl)propyl]thio]phenol (5)

5

10



15

A solution of 3 (1.08 g, 3.0 mmol) and imidazo[4,5-c]pyridine 4 (538 mg, 4.5 mmol) in dimethylformamide (30 ml) was heated at 70°C. After 20 hours, the solvent was removed and the crude residue (1.54 g) was chromatographed (silica gel; methylene chloride/methanol/ammonium hydroxide 90/10/1) to give 5 (745 mg, 62%) as a crystalline solid. mp (DSC) 181°C. The structural assignment was supported by the <sup>1</sup>H-NMR spectrum. Analysis calculated for C<sub>23</sub>H<sub>31</sub>N<sub>3</sub>OS·0.75 H<sub>2</sub>O:


25 C, 67.20; H, 7.97; N, 10.22; S, 7.80. Found: C, 66.85; H, 7.67; N, 10.08; S, 7.44.

- 38 -

Example 42,6-bis(1,1-dimethylethyl)-4-[(2-hydroxyethyl)-  
thiol]phenol (6)

5

10



15 To a stirred mixture of  $K_2CO_3$  (52.2 g, 0.378 mol) in a solution of 1 (100.0 g, 0.373 mol) in DMF (900 ml) at room temperature was added bromoethanol (28.2 ml, 0.378 mol). After 6 hours, the reaction was recharged with an additional 6 ml of bromoethanol and stirred for

20 one additional hour. The reaction was concentrated in vacuo, poured onto water, and extracted several times with ethyl acetate. The combined ethyl acetate layers were dried ( $Na_2SO_4$ ), filtered and evaporated to yield a brown oil. The crude product was chromatographed on

25 silica gel using mixtures of ethyl acetate and hexane as eluents. The oil thus obtained was crystallized under hexane to give 83.93 g (78.7%) of crystalline product in two crops, mp 62.6-65°. An analytical sample was obtained by recrystallization from hexane.

30 The structural assignment was supported by the  $^1H$  NMR spectrum. Analysis calculated for  $C_{16}H_{26}O_2S$ : C, 68.04; H, 9.28; S, 11.35. Found: C, 68.06; H, 9.46; S, 11.44.




- 39 -

Example 54-[(2-bromoethyl)thio]-2,6-bis(1,1-dimethylethyl)phenol (7)

5

10




Triphenylphosphine (3.72 g, 14.2 mmol) was added  
15 to a stirred solution of **6** (2.0 g, 7.1 mmol) and carbon  
tetrabromide (4.7 g, 14.2 mmol) in dry ether (80 ml).  
After stirring at room temperature for 16 hours, the  
reaction mixture was concentrated and the crude product  
was chromatographed (silica gel; hexane/ethyl acetate  
20 95/5) to give **7** (2.1 g, 82%) as a crystalline solid.  
mp (DSC) 112°C. Analysis calculated for  $C_{16}H_{25}BrOS$ :  
C, 55.65; H, 7.30; Br, 23.14; S, 9.28. Found C, 55.49;  
H, 7.27; Br, 22.10; S, 9.36.

- 40 -

Example 6

2,6-bis(1,1-dimethylethyl)-4-[[2-(5H-imidazo-  
[4,5-c]pyridin-5-yl)ethyl]thio]-  
5 phenol hydrate (8)


10



15

A solution of 7 (1.1 g, 3.9 mmol) and imidazo[4,5-c]pyridine 4 (464 mg, 3.9 mmol) in dimethylacetamide (40 ml) was heated at 70°C. After 48 hours, the solvent was removed and the crude residue (1.2 g) was  
20 chromatographed (silica gel; methylene chloride/methanol/ammonium hydroxide 90/10/1) to give 8 (252 mg, 17%) as a crystalline solid. mp (DSC) 196°C. Analysis calculated for  $C_{27}H_{29}N_3OS \cdot 0.25 H_2O$ : C, 68.09; H, 7.66; N, 10.86; S, 8.26. Found: C, 68.17; H, 7.93;  
25 N, 10.66; S, 8.23.

- 41 -

Example 72,6-bis(1,1-dimethylethyl)-4-[[3-(1,3-dioxolan-2-yl)propyl]thio]phenol (10)


To a solution of 1 (15g, 63 mmol) in acetone (300 ml), anhydrous potassium carbonate (17.4 g, 126 mmol) and 2-(3-chloropropyl)-1,3-dioxalane 9 (9.9 ml) were added. After stirring at room temperature for 48 hours, the reaction mixture was filtered and the filter cake was washed with more acetone. The combined filtrates were concentrated and the crude black liquid (25.7g) was chromatographed (silica gel; hexane/ethyl acetate 9/1) to give 13.8 g of a mixture of product 10 and starting material 9 (7:3). This mixture was used in the next example without further purification.

- 42 -

Example 84-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]thio]butanal (11)

5

10




A mixture of crude 10 (13.1 g) as obtained in  
15 Example 7 was heated in acetic acid (180 ml) and water  
(120 ml) at 70-75°C. After 4 hours, the reaction  
mixture was cooled to room temperature and then poured  
onto ether. The ethereal layer was separated and  
washed with water, an aqueous sodium bicarbonate  
20 solution and brine. After drying over MgSO<sub>4</sub>, the  
ethereal fractions were filtered and concentrated. The  
crude residue (11.8 g) was chromatographed (silica gel:  
hexane/ethyl acetate 9/1) to give 11 (6.1 g) as a  
colorless liquid. Analysis calculated for  
25 C<sub>18</sub>H<sub>28</sub>O<sub>2</sub>S·0.4.H<sub>2</sub>O: C, 68.42; H, 9.20; S, 10.16. Found: C,  
68.42; H, 8.76; S, 10.65.

- 43 -

Example 92,6-bis(1,1-dimethylethyl)-4-[(4-hydroxybutyl)-  
thiolphenol (12)

5

10




To a solution of 11 (5.72 g, 18.57 mmol) in  
15 ethanol (20 ml), sodium borohydride (2.1 g, 55.7 mmol)  
was added and the mixture was refluxed for 5 hours.  
The reaction mixture was cooled to room temperature and  
then quenched by adding an excess of water. After  
extraction in methylene chloride (2 X 500), the organic  
20 layer was washed (with brine), dried (over  $\text{MgSO}_4$ ) and  
filtered. The solvent was removed from the organic  
fractions and the crude residue (6.2 g) was  
chromatographed (silica gel; hexane/ethyl acetate  
75/25) to give 12 (4.52 g, 78%) as a colorless liquid.  
25 The structural assignment was supported by the  $^1\text{H}$ -NMR  
spectrum. Analysis calculated for  $\text{C}_{18}\text{H}_{30}\text{O}_2\text{S} \cdot 0.25\text{H}_2\text{O}$ : C,  
68.63; H, 9.76; S, 10.18. Found: C, 68.50; H, 9.48;  
S, 10.28.

- 44 -

Example 104-[(4-bromobutyl)thio]-2,6-bis(1,1-dimethylethyl)phenol (13)

5

10




Triphenylphosphine (1.69 g, 6.45 mmol) was added  
15 to a stirred solution of **12** (1.0 g, 3.22 mmol) and  
carbon tetrabromide (2.14 g, 6.45 mmol) in dry ether  
(50 ml). After stirring at room temperature for 18  
hours, the reaction mixture was concentrated and the  
crude product chromatographed (silica gel; hexane/ethyl  
20 acetate 90/10) to give **13**. The structural assignment  
was supported by the <sup>1</sup>H-NMR spectrum.

- 45 -

Example 11

2,6-bis(1,1-dimethylethyl)-4-[[4-(5H-imidazo-  
[4,5-c]pyridin-5-yl)butyl]thio-  
5 phenol hydrate (14)



A solution of 13 (2.0 g, 5.36 mmol) and  
imidazo[4,5-c]pyridine 4 (640 mg, 5.36 mmol) in  
dimethylacetamide (40 ml) was heated at 70°C. After 48  
hours, the solvent was removed. The crude residue was  
redissolved in methylene chloride and washed with  
20 aqueous potassium carbonate and brine. The organic  
fractions were dried over MgSO<sub>4</sub>, filtered and  
concentrated. The residue obtained (2.9 g) was  
chromatographed (silica gel; methylene  
chloride/methanol/ammonium hydroxide 90/10/1) to give  
25 14 (510 mg, 23%). Analysis calculated for  
C<sub>24</sub>H<sub>33</sub>N<sub>3</sub>OS·0.25H<sub>2</sub>O: C, 69.28; H, 8.11; N, 10.10; S, 7.71.  
Found: C, 69.05; H, 8.22; N, 9.84; S, 7.80.

- 46 -

Example 12

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-  
imidazo[5,4-c]pyridin-1-yl)propyl]-  
thiolphenol (15)


5

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-  
imidazo[4,5-c]pyridin-1-yl)propyl]-  
thiolphenol (16)

10


2,6-bis(1,1-dimethylethyl)-4-[[3-(5H-imidazo-  
[4,5-c]pyridin-5-yl)propyl]thiolphenol (5)

15




20

25



30





- 47 -

To a solution of imidazopyridine 4 (5 g, 42 mmol) in dimethylformamide (250 ml), sodium hydride (2.01 g, 60% dispersion in mineral oil, 50.4 mmol) was added. After stirring for 1 hour at 20°C, 1-chloro-3-iodopropane (5.42 ml, 50.4 mmol) was added over 10 minutes. The reaction mixture was stirred for 24 hours and then the phenol 1 (14 g, 58 mmol) was added over 5 minutes. After stirring for an additional 48 hours, the reaction mixture was poured over ice and extracted with methylene chloride. The organic layer was separated, washed (brine), dried (MgSO<sub>4</sub>), filtered and concentrated. The crude product was chromatographed (silica gel; methylene chloride/methanol/ammonium hydroxide 95/5/0.5) to give 15, 16 and 5, respectively. The structural assignments were supported by the <sup>1</sup>H-NMR spectrum.

15: mp (DSC) 132°C; Analysis calculated for C<sub>23</sub>H<sub>31</sub>N<sub>3</sub>OS: C, 69.48; H, 7.86; N, 10.57; S, 8.06. Found C, 69.16; H, 7.72; N, 10.31; S, 8.03.

16: mp (DSC) 131°C; Analysis calculated for C<sub>23</sub>H<sub>31</sub>N<sub>3</sub>OS: C, 69.48; H, 7.86; N, 10.57; S, 8.06. Found C, 69.17; H, 7.91; N, 10.53; S, 7.96.



25

- 48 -

Example 13

(2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-  
imidazo[5,4-b]pyridin-1-yl)propyl]thio]-  
phenol hydrate (18)

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-  
imidazo[4,5-b]pyridin-1-yl)propyl]thio]-  
phenol hydrate (19)



To a solution of imidazopyridine 17 (5 g, 42 mmol) in dimethylformamide (250 ml) was added sodium hydride (2.01 g, 60% dispersion in mineral oil, 50.4 mmol). After stirring for 1 hour at 20°C, 1-chloro-3-iodopropane (5.42 ml, 50.4 mmol) was added over 10 minutes. The reaction mixture was stirred for 24 hours and then the phenol 1 (12.8 g, 54 mmol) was added over 5 minutes. After stirring for an additional 48 hours, the reaction mixture was poured over ice and extracted with methylene chloride. The organic layer was

- 49 -

separated, washed (brine), dried ( $\text{MgSO}_4$ ), filtered and concentrated. The crude product (24.8 g) was chromatographed (silica gel; methylene chloride/methanol/ammonium hydroxide 95/5/0.5) to give  
5 18 & 19 respectively. The structural assignments were supported by the  $^1\text{H}$ -NMR spectrum.

18: mp (DSC)  $97^\circ\text{C}$ ; Analysis calculated for  $\text{C}_{23}\text{H}_{31}\text{N}_3\text{OS} \cdot 0.25\text{H}_2\text{O}$ : C, 68.70; H, 7.90; N, 10.45; S, 7.97. Found:  
10 C, 68.58; H, 7.70; N, 10.28; S, 7.95.


19: mp (DSC)  $113^\circ\text{C}$ ; Analysis calculated for  $\text{C}_{23}\text{H}_{31}\text{N}_3\text{OS} \cdot 0.75\text{H}_2\text{O}$ : C, 67.20; H, 7.97; N, 10.22; S, 7.80. Found:  
15 C, 67.53; H, 7.72; N, 9.95; S, 7.68.

- 50 -

Example 142,6-bis(1,1-dimethylethyl)-4-[[3-(1H-benzimidazol-1-yl)propyl]thio]phenol (21)

5

10




15 To a solution of benzimidazole 20 (5 g, 42 mmol) in dimethylformamide (200 ml) was added sodium hydride (2.01 g, 60% dispersion in mineral oil, 50.4 mmol). After stirring for 1 hour at 20°C, 1-chloro-3-iodopropane (5.42 ml, 50.4 mmol) was added over 10  
20 minutes. The reaction mixture was stirred for 24 hours and then the phenol 1 (14 g, 58 mmol) and triethylamine (7 ml) were added over 5 minutes. After stirring for an additional 48 hours, the reaction mixture was poured over ice and extracted with methylene chloride. The  
25 organic layer was separated, washed (brine), dried (MgSO<sub>4</sub>), filtered and concentrated. The crude product (31 g) was chromatographed (silica gel; methylene chloride/methanol/ammonium hydroxide 95/5/0.5) to give 21 (7.43 g, 45%). mp (DSC) 96°C. Analysis calculated  
30 for C<sub>24</sub>H<sub>32</sub>N<sub>2</sub>OS: C, 72.68; H, 8.13; N, 7.06; S, 8.08. Found: C, 72.51; H, 8.18; N, 7.02; S, 8.06.

- 51 -

Example 151-(3-chloropropyl)-1H-indole (23)

5



10


To a solution of indole 22 (5 g, 42 mmol) in  
15 dimethylformamide (200 ml) was added sodium hydride  
(2.01 g, 60% dispersion in mineral oil, 50.4 mmol).  
After stirring for 1 hour at 20°C, 1-chloro-3-  
iodopropane (5.42 ml, 50.4 mmol) was added over 10  
minutes. The reaction mixture was stirred for 24 hours  
20 and then poured over ice and extracted with methylene  
chloride. The organic layer was separated, washed  
(brine), dried (MgSO<sub>4</sub>), filtered and concentrated. The  
crude product (25.3 g) was chromatographed (silica gel;  
hexane/ethyl acetate 95/5) to give 23 (4.02 g, 49%).  
25 The structural assignment was supported by the <sup>1</sup>H-NMR  
spectra in CDCl<sub>3</sub>.

- 52 -

Example 162,6-bis(1,1-dimethylethyl)-4-[[3-(1H-indol-1-yl)propyl]thio]phenol hydrate (24)

5

10



15

To a solution of **23** (4.02 g, 20.7 mmol) and **1** (4.92 g, 20.7 mmol) in dimethylformamide (100 ml), triethylamine (5.76 ml, 41.36 mmol) was added. After stirring for 72 hours, the reaction mixture was poured over ice and extracted with methylene chloride. The organic layer was separated, washed (brine), dried (MgSO<sub>4</sub>), filtered and concentrated. The crude product (8.44 g) was chromatographed (silica gel; hexane/ethyl acetate 95/5) to give **24** (2.86 g, 34%) as a colorless liquid. The structural assignment was supported by the

25


<sup>1</sup>H-NMR spectrum. Analysis calculated for C<sub>25</sub>H<sub>33</sub>NOS·0.25H<sub>2</sub>O: C, 75.05; H, 8.44; N, 3.50; S, 8.01. Found: C, 74.88; H, 8.32; N, 3.47; S, 7.69.

- 53 -

Example 172,6-bis(1,1-dimethylethyl)-4-[[2-[(1,3-benzoxazol-2-yl)thio]ethyl]thio]phenol hydrate (26)

5

10



15 To a solution of 2-mercaptobenzoxazole 25 (460 mg, 3 mmol) in dimethylformamide (30 ml), silver tetrafluoroborate (584 mg, 3 mmol) was added. After 5 minutes, 7 (1.04 g, 3 mmol) was added and the mixture continued to stir under argon at room temperature.

20 After 18 hours, the reaction mixture was filtered and the residue was washed with methylene chloride. The combined filtrates were concentrated and the crude yellowish liquid (4 g) obtained was chromatographed (silica gel; hexane/ethyl acetate 95/5) to give 26 (210

25 mg, 17%) as a crystalline solid. mp (DSC) 118°C. Analysis calculated for  $C_{23}H_{29}NO_2S_2 \cdot 0.4H_2O$ : C, 65.34; H, 7.10; N, 3.31; S, 15.17. Found: C, 65.38; H, 6.70; N, 3.22; S, 14.98.


- 54 -

Example 18

(±) 2,6-bis(1,1-dimethylethyl)-4-[[2-hydroxy-  
3-(5H-imidazo[4,5-c]pyridin-5-yl)propyl-  
thio]phenol hydrate (27)

5

10



15

A solution of 2,6-di-t-butyl-4-(2'-hydroxy-3'-chloropropyl)thiophenol (3 g, 9 mmol) and 1(H)-imidazo(4,5c)pyridine (2.1 g, 18 mmol) in DMA (30 mls) was heated to 70°C for 14 hours under N<sub>2</sub>. The solvent was removed via high vacuum and the residue was treated with CH<sub>2</sub>Cl<sub>2</sub> and H<sub>2</sub>O. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>) and solvent removed. The residue was chromatographed with 15 MeOH/85 CH<sub>2</sub>Cl<sub>2</sub>/.5 NH<sub>4</sub>OH. The product 27 (1.6g, 43%) had a melting point of 185-187°C. An analytical sample was obtained by recrystallization from MeOH/H<sub>2</sub>O.

20

25

C<sub>23</sub>H<sub>31</sub>N<sub>3</sub>O<sub>2</sub>S.1/2 H<sub>2</sub>O M.W.=422.59. Analysis Calculated for C<sub>23</sub>H<sub>31</sub>N<sub>3</sub>O<sub>2</sub>S.1/2 H<sub>2</sub>O: C, 65.37; H, 7.63; N, 9.94; S, 7.59. Found: C, 64.94; H, 8.05; N, 9.61; S, 7.59.




- 55 -


Example 196-[[1,3-bis(dimethylethyl)-4-hydroxy phenyl]-  
thiomethyl]-1-methyl-1H-guinolin-2-one (28)

5

10

(a) 1,6-Dimethylquinoline methylsulfate (28a):

15




20

A solution of 6-methylquinoline (39 g, 0.272 mol) and dimethyl sulfate (30 mls, 0.29 mol) in toluene (400 ml) was stirred rapidly using mechanical stirring at 60°C. The resulting thick paste was diluted with more toluene, cooled to 25°C and filtered. The product (60 g, 82%) was dried in a vacuum oven at 40°C overnight. As this material was very hygroscopic, it was stored in a dessicator over P<sub>2</sub>O<sub>5</sub>.

25

(b) 1,6-Dimethyl-guinolin-2-one (28b)

30




35

A solution of the salt 28a (60 g, 0.22 moles) and potassium ferricyanide (147 g, 0.45 mol) in water

- 56 -

(300 ml) was stirred rapidly as a solution of NaOH (27 g, 0.68 mol) in water (100 mls) was added slowly over 45 minutes, maintaining a reaction temperature of 50°C. After cooling, the mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>, dried (MgSO<sub>4</sub>) and evaporated. The resulting crude solid was recrystallized from benzene to give pure title compound **28b** (23.7 g, 62%). The structural assignment was supported by the <sup>1</sup>H-NMR spectrum. Analysis Calculated for C<sub>11</sub>H<sub>11</sub>NO: C, 76.28; H, 6.40; N, 8.09. Found: C, 76.31; H, 6.45; N, 7.99.


(c) 6-Bromomethyl-1-methylquinoline-2-one (28c)



A stirred solution of **28b** (22 g, 0.127 mol) and N-bromosuccinimide (23 g, 0.13 mol) in CCl<sub>4</sub> (400 mls) was irradiated for 2.5 hours with a 275 watt heat lamp after an addition of AIBN (50 mg) to initiate the free radical reaction. During this time, the reaction achieved reflux. After cooling, the mixture was concentrated to a small volume and ethyl acetate (400 mls) and water (150 mls) were added and the layers separated. The organic phase was dried (MgSO<sub>4</sub>) and evaporated. The crude material was purified by chromatography on silica gel using 35% ethyl acetate/toluene to give the product **28c** (10.4 g, 33%) as a solid, mp 99-100°C. The structural assignment was supported by the <sup>1</sup>H-NMR spectrum. Analysis calculated for C<sub>11</sub>H<sub>10</sub>NOBr: C, 52.41; H, 4.00; N, 5.56; Br, 31.69. Found: C, 52.11; H, 3.91; N, 5.40; Br, 31.40.

- 57 -

(d) 6-[[1,3-bis(dimethylethyl)-4-hydroxy phenyl]-thiomethyl]-1-methyl-1H-quinolin-2-one (28)



10 A mixture of K<sub>2</sub>CO<sub>3</sub> (850 mg, 6.2 mmol) in a solution  
of **28c** (750 mg, 0.3 mmol) and 2,6-di-*t*-butyl-4-  
mercaptophenol (1 g, 3 mmol) in DMF (25 ml) was stirred  
for 4 hours at 25°C. The mixture was filtered and the  
filtrate was evaporated in vacuo using an oil pump.  
15 The residue was dissolved in ethyl acetate, washed with  
water and the organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>). After  
evaporation, the residue was chromatographed on silica  
gel with 40% ethyl acetate/hexane to give pure product  
(950 mg, 77%) as a solid, mp 135-136°C. The structural  
20 assignment was supported by the <sup>1</sup>H-NMR spectrum.  
Analysis calculated for C<sub>25</sub>H<sub>31</sub>NO<sub>2</sub>S: C, 73.31; H, 7.63; N,  
3.42; S, 7.83. Found: C, 73.28; H, 7.52; N, 3.36; S,  
7.57.

25


- 58 -

Example 20

2,6-bis(1,1-dimethylethyl)-4-[[ (3H-imidazo-  
[4,5-b]pyridin-2-yl)methyl]thio]phenol (29)


5

10



(a) 2-Chloromethyl-1(H)-imidazo[4,5-b]pyridine (29a)

15




20

To a slurry of 2-hydroxymethyl-1(H)-imidazo[4,5-b]pyridine (2 g, 13 mmol) in EtOH (5 mls) was added a solution of HCl in dioxane (4M in HCl, 10 ml, 40 mmol). This was stirred for 10 minutes, and the crude hydrochloride salt was filtered. This salt was slurried in CHCl<sub>3</sub> (15 ml), thionyl chloride (5 ml) was added and the mixture was refluxed for 48 hours. The reaction mixture was filtered hot and evaporated. The crude product was used without further purification.

30

- 59 -






(b) 2,6-bis(1,1-dimethylethyl)-4-[[[3H-imidazo-  
[4,5-b]pyridin-2-yl)methyl]thio]phenol (29)



10

A mixture of  $K_2CO_3$  (10 g), crude **29a** (2.6 g, 12 mmol) and 2,6-di-*t*-butyl-4-mercaptophenol (5 g, 21 mmol) in DMF (50 mls) was stirred for 16 hours at 25°C. The reaction was filtered and the filtrate was concentrated in vacuo using an oil pump. The residue was chromatographed on silica gel using MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (5:94:1) as the eluent to give the purified product **29** (3 g, 80%) as a crystalline solid, mp 241-242°C. The structural assignment was supported by the <sup>1</sup>H-NMR spectrum. Analysis calculated for C<sub>21</sub>H<sub>27</sub>N<sub>3</sub>OS: C, 68.26; H, 7.36; N, 11.37; S, 8.68. Found: C, 67.91; H, 7.44; N, 10.97; S, 8.69.

- 60 -

Compound No.	Compound Structure
1	 <chem>CC(C)(C)S[C@@H]1C=C(C(C)(C)O)C(C(C)(C)O)C1C(C)(C)C</chem>
2	 <chem>CC(C)(C)S[C@@H]1C=C(C(C)(C)O)C(C(C)(C)O)C1C(C)(C)CCO</chem>
3	 <chem>CC(C)(C)S[C@@H]1C=C(C(C)(C)O)C(C(C)(C)O)C1C(C)(C)CCBr</chem>
4	 <chem>C1=NC2=C(N1)N=CN=C2</chem>
5	 <chem>CC(C)(C)S[C@@H]1C=C(C(C)(C)O)C(C(C)(C)O)C1C(C)(C)CCN2C=NC3=C(N2)N=CN=C3</chem>






5






10

15






20







25

Compound No.	Compound Structure
6	
7	
8	
9	
10	

Compound No.	Compound Structure
11	
12	
13	
14	
15	



Compound No.	Compound Structure
5       <b>16</b>	
10       <b>17</b>	
15       <b>18</b>	
20       <b>19</b>	
25       <b>20</b>	

Compound No.	Compound Structure
21	
22	
23	
24	
25	
26	






5

10



15

20

25

Compound No.	Compound Structure
27	
28	
28a	
28b	
28c	

- 66 -

Compound No.	Compound Structure
29	
29a	

- 67 -


The foregoing examples are provided to enable one of ordinary skill in the art to practice the present invention. These examples are merely illustrative, however, and should not be read as limiting the scope of the invention as it is claimed in the appended claims.

While the present invention has been described herein with some specificity, and with reference to certain preferred embodiments thereof, those of ordinary skill in the art will recognize numerous variations, modifications and substitutions of that which has been described which can be made, and which are within the scope and spirit of the invention. For example, effective dosages other than the preferred ranges set forth hereinabove may be applicable as a consequence of variations in the responsiveness of the animal being treated, dosage-related adverse effects, if any, and analogous considerations. Likewise, the specific pharmacological responses observed may vary according to, and depending upon, the particular active compound selected, or whether there are present certain pharmaceutical carriers, as well as the type of formulation and mode of administration employed. Such expected variations and/or differences in the results are contemplated in accordance with the objects and practices of the present invention. It is intended therefore that all of these modifications and variations be within the scope of the present invention as described and claimed herein, and that the invention be limited only by the scope of the claims which follow, and that such claims be interpreted as broadly as is reasonable.

- 68 -

What is claimed is:

1. A compound of the formula:



or a pharmaceutically acceptable salt thereof, wherein:

$R^1$  and  $R^2$  are each alkyl;

$n$  is an integer of from 1 to 5;

$p$  is an integer of from 0 to 1;

$q$  is an integer of from 0 to 3;


$r$  is an integer of from 0 to 1; and

$R^3$  is a bicyclic group,


with the proviso that when  $R^1$  and  $R^2$  are each alkyl, and  $p$ ,  $q$  and  $r$  are each 0,  $R^3$  is not a naphthylmethyl group.

- 69 -


2. A compound of Claim 1 having the structure:




3. A compound of Claim 1 having the structure:



4. A compound of Claim 1 having the structure:




5. A compound of Claim 1 having the structure:




- 70 -


6. A compound of Claim 1 having the structure:



7. A compound of Claim 1 having the structure:




8. A compound of Claim 1 having the structure:






- 71 -


9. A compound of Claim 1 having the structure:



10. A compound of Claim 1 having the structure:




11. A compound of Claim 1 having the structure:




- 72 -


12. A compound of Claim 1 having the structure:



13. A compound of Claim 1 having the structure:




14. A compound of Claim 1 having the structure:



15. A pharmaceutical composition comprising a pharmaceutically-acceptable carrier and a pharmaceutically-effective amount of a compound of the formula:

- 73 -



or a pharmaceutically acceptable salt thereof, wherein:

$\text{R}^1$  and  $\text{R}^2$  are each alkyl;

$n$  is an integer of from 1 to 5;

$p$  is an integer of from 0 to 1;

$q$  is an integer of from 0 to 3;

$r$  is an integer of from 0 to 1; and

$\text{R}^3$  is a bicyclic group,

with the proviso that when  $\text{R}^1$  and  $\text{R}^2$  are each alkyl, and  $p$ ,  $q$  and  $r$  are each 0,  $\text{R}^3$  is not a naphthylmethyl group.

16. The pharmaceutical composition of Claim 15 wherein the compound is:

2,6-bis(1,1-dimethylethyl)-4-[[3-(5H-imidazo-  
[4,5-c]pyridin-5-yl)propyl]thio]phenol (5);

2,6-bis(1,1-dimethylethyl)-4-[[2-(5H-imidazo-  
[4,5-c]pyridin-5-yl)ethyl]thio]-  
phenol hydrate (8);

2,6-bis(1,1-dimethylethyl)-4-[[4-(5H-imidazo-  
[4,5-c]pyridin-5-yl)butyl]thio-  
phenol hydrate (14);

- 74 -

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-imidazo[5,4-c]pyridin-1-yl)propyl]-thio]phenol (15);

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-imidazo[4,5-c]pyridin-1-yl)propyl]-thio]phenol (16);

(2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-imidazo[5,4-b]pyridin-1-yl)propyl]thio]-phenol hydrate (18);

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-imidazo[4,5-b]pyridin-1-yl)propyl]thio]-phenol hydrate (19);

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-benzimidazol-1-yl)propyl]thio]phenol (21);

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-indol-1-yl)propyl]thio]phenol hydrate (24);

2,6-bis(1,1-dimethylethyl)-4-[[2-[(1,3-benzoxazol-2-yl)thio]ethyl]thio]phenol hydrate (26);


(±) 2,6-bis(1,1-dimethylethyl)-4-[[2-hydroxy-3-(5H-imidazo[4,5-c]pyridin-5-yl)propyl]-thio]phenol hydrate (27);

6-[[1,3-bis(dimethylethyl)-4-hydroxy phenyl]-thiomethyl]-1-methyl-1H-quinolin-2-one (28); or

2,6-bis(1,1-dimethylethyl)-4-[[3-(3H-imidazo[4,5-b]pyridin-2-yl)methyl]thio]phenol (29).

- 75 -

17. A method for treating 5-lipoxygenase mediated conditions in an animal in need of such treatment comprising administering to the animal a therapeutically-effective amount of a compound of the formula:



or a pharmaceutically acceptable salt thereof, wherein:

$\text{R}^1$  and  $\text{R}^2$  are each alkyl;

$n$  is an integer of from 1 to 5;

$p$  is an integer of from 0 to 1;

$q$  is an integer of from 0 to 3;

$r$  is an integer of from 0 to 1; and

$\text{R}^3$  is a bicyclic group,

with the proviso that when  $\text{R}^1$  and  $\text{R}^2$  are each alkyl, and  $p$ ,  $q$  and  $r$  are each 0,  $\text{R}^3$  is not a naphthylmethyl group.

18. The method of Claim 17 wherein the compound is:

2,6-bis(1,1-dimethylethyl)-4-[[3-(5H-imidazo-  
[4,5-c]pyridin-5-yl)propyl]thio]phenol (5);

2,6-bis(1,1-dimethylethyl)-4-[[2-(5H-imidazo-  
[4,5-c]pyridin-5-yl)ethyl]thio]-  
phenol hydrate (8);

- 76 -

2,6-bis(1,1-dimethylethyl)-4-[[4-(5H-imidazo-  
[4,5-c]pyridin-5-yl)butyl]thio-  
phenol hydrate (14);

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-  
imidazo[5,4-c]pyridin-1-yl)propyl]-  
thio]phenol (15);

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-  
imidazo[4,5-c]pyridin-1-yl)propyl]-  
thio]phenol (16);

(2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-  
imidazo[5,4-b]pyridin-1-yl)propyl]thio]-  
phenol hydrate (18);

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-  
imidazo[4,5-b]pyridin-1-yl)propyl]thio]-  
phenol hydrate (19);

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-benzimidazol-  
1-yl)propyl]thio]phenol (21);

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-indol-  
1-yl)propyl]thio]phenol hydrate (24);

2,6-bis(1,1-dimethylethyl)-4-[[2-[(1,3-benzoxazol-  
2-yl)thio]ethyl]thio]phenol hydrate (26);


(±) 2,6-bis(1,1-dimethylethyl)-4-[[2-hydroxy-  
3-(5H-imidazo[4,5-c]pyridin-5-yl)propyl-  
thio]phenol hydrate (27);

6-[[1,3-bis(dimethylethyl)-4-hydroxy phenyl]-  
thiomethyl]-1-methyl-1H-quinolin-2-one (28); or

- 77 -

2,6-bis(1,1-dimethylethyl)-4-[[ (3H-imidazo-  
[4,5-b]pyridin-2-yl)methyl]thio]phenol (29).

19. A method for treating an inflammation-associated disorder or an allergy in an animal in need of such treatment comprising administering to the animal a therapeutically-effective amount of a compound of the formula:



or a pharmaceutically acceptable salt thereof, wherein:

$\text{R}^1$  and  $\text{R}^2$  are each alkyl;

$n$  is an integer of from 1 to 5;

$p$  is an integer of from 0 to 1;

$q$  is an integer of from 0 to 3;

$r$  is an integer of from 0 to 1; and

$\text{R}^3$  is a bicyclic group,

with the proviso that when  $\text{R}^1$  and  $\text{R}^2$  are each alkyl, and  $p$ ,  $q$  and  $r$  are each 0,  $\text{R}^3$  is not a naphthylmethyl group.

20. The compound of Claim 19 wherein the compound is:

2,6-bis(1,1-dimethylethyl)-4-[[3-(5H-imidazo-  
[4,5-c]pyridin-5-yl)propyl]thio]phenol (5);

- 78 -

2,6-bis(1,1-dimethylethyl)-4-[[2-(5H-imidazo-  
[4,5-c]pyridin-5-yl)ethyl]thio]-  
phenol hydrate (8);

2,6-bis(1,1-dimethylethyl)-4-[[4-(5H-imidazo-  
[4,5-c]pyridin-5-yl)butyl]thio]-  
phenol hydrate (14);

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-  
imidazo[5,4-c]pyridin-1-yl)propyl]-  
thio]phenol (15);

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-  
imidazo[4,5-c]pyridin-1-yl)propyl]-  
thio]phenol (16);

(2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-  
imidazo[5,4-b]pyridin-1-yl)propyl]thio]-  
phenol hydrate (18);

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-  
imidazo[4,5-b]pyridin-1-yl)propyl]thio]-  
phenol hydrate (19);

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-benzimidazol-  
1-yl)propyl]thio]phenol (21);

2,6-bis(1,1-dimethylethyl)-4-[[3-(1H-indol-  
1-yl)propyl]thio]phenol hydrate (24);

2,6-bis(1,1-dimethylethyl)-4-[[2-[(1,3-benzoxazol-  
2-yl)thio]ethyl]thio]phenol hydrate (26);

(±) 2,6-bis(1,1-dimethylethyl)-4-[[2-hydroxy-  
3-(5H-imidazo[4,5-c]pyridin-5-yl)propyl-  
thio]phenol hydrate (27);



- 79 -

6-[[1,3-bis(dimethylethyl)-4-hydroxy phenyl]-  
thiomethyl]-1-methyl-1H-quinolin-2-one (28); or

2,6-bis(1,1-dimethylethyl)-4-[[ (3H-imidazo-  
[4,5-b]pyridin-2-yl)methyl]thio]phenol (29).

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 95/12263

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C07D215/22 C07D209/08 C07D263/58 C07D235/06 C07D471/04  
A61K31/44 A61K31/415 A61K31/42 //(C07D471/04, 235:00,  
221:00)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 190 684 (G. D. SEARLE) 13 August 1986 see claims 1,16 ---	1,17,19
X	CHEMICAL ABSTRACTS, vol. 116, no. 25, 22 June 1992, Columbus, Ohio, US; abstract no. 255437, S. A. ZOTOVA ET AL 'Synthesis and biological activity of substituted sulfides of indoles and benzofuran' page 771 ; see abstract & KHIM.-FARM. ZH., vol.26, no.1, 1992 pages 52 - 55 --- -/--	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

23 January 1996

Date of mailing of the international search report

-7.02.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Voyiazoglou, D

# INTERNATIONAL SEARCH REPORT

Inter. Application No.  
PCT/US 95/12263

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 479 631 (ADIR ET COMPANIE) 8 April 1992 see claim 1 ---	1
X	DE,A,29 31 741 (DR. KARL THOMAE) 26 February 1981 see claim 1 ---	1
X	DE,A,31 43 327 (DR. KARL THOMAE) 9 June 1982 see claim 1 ---	1
P,X	CHEMICAL ABSTRACTS, vol. 122, no. 15, 10 April 1995, Columbus, Ohio, US; abstract no. 187387, KOJI KOBAYASHI ET AL 'Preparation of phthalimide derivatives as phospholipase A2 inhibitors' page 1037 ; see abstract & JP,A,6 312 978 (NIPPON TOBACCO SANGYO & YOSHITOMI) 8 November 1994 ---	1,17,19
A	EP,A,0 417 745 (G. D. SEARLE & CO) 20 March 1991 see page 3; claim 1 -----	1,19

## INTERNATIONAL SEARCH REPORT

1. International application No.

PCT/US95/12263

**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Although claims 17-19 are directed to a method of treatment of (diagnostic method practised on) the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

**Remark on Protest**

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 95/12263

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0190684	13-08-86	AU-B- 589942	26-10-89
		AU-B- 5288386	07-08-86
		CA-A- 1332057	20-09-94
		JP-B- 6062549	17-08-94
		JP-A- 61191670	26-08-86
		US-A- 5019597	28-05-91
		US-A- 5157053	20-10-92
		US-A- 5198435	30-03-93
		US-A- 5244899	14-09-93
EP-A-0479631	08-04-92	FR-A- 2666583	13-03-92
		AT-T- 113595	15-11-94
		AU-B- 638354	24-06-93
		AU-B- 8369491	12-03-92
		CA-A- 2050730	07-03-92
		DE-D- 69104948	08-12-94
		DE-T- 69104948	08-06-95
		ES-T- 2066390	01-03-95
		JP-A- 4247082	03-09-92
		JP-B- 7086106	20-09-95
		US-A- 5206247	27-04-93
DE-A-2931741	26-02-81	AU-B- 539070	13-09-84
		AU-B- 6100680	12-02-81
		BE-A- 884628	04-02-81
		JP-A- 56036452	09-04-81
		US-A- 4329347	11-05-82
DE-A-3143327	09-06-82	AT-B- 376661	27-12-84
		AT-B- 375923	25-09-84
		AU-B- 547929	14-11-85
		AU-B- 7739081	20-05-82
		BE-A- 891103	12-05-82
		CA-A- 1183848	12-03-85
		CH-A- 652120	31-10-85
		FR-A, B 2493841	14-05-82
		GB-A, B 2098595	24-11-82
		JP-A- 57114569	16-07-82
		LU-A- 83744	13-04-83
		NL-A- 8105100	01-06-82

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 95/12263

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE-A-3143327		SE-B- 452156 SE-A- 8106704 SU-A- 1107759 US-A- 4629733	16-11-87 13-05-82 07-08-84 16-12-86
JP-A-6312978	08-11-94	NONE	
EP-A-0417745	20-03-91	US-A- 4990518 CA-A- 2025084 DE-D- 69004389 DE-T- 69004389 ES-T- 2059942 JP-A- 3109386	05-02-91 14-03-91 09-12-93 24-03-94 16-11-94 09-05-91